

Sample Coding

TYPE	CODE	DESCRIPTION
Diagnosis: ICD-10-CM	C50.011–C50.019 C50.111–C50.119 C50.211–C50.219 C50.311–C50.319 C50.411–C50.419 C50.511–C50.519 C50.611–C50.619 C50.811–C50.819 C50.911–C50.919	Malignant neoplasm of the female breast
	C50.021–C50.029 C50.121–C50.129 C50.221–C50.229 C50.321–C50.329 C50.421–C50.429 C50.521–C50.529 C50.621–C50.629 C50.821–C50.829 C50.921–C50.929	Malignant neoplasm of the male breast
Drug: HCPCS code	J3490	Not otherwise classified drugs
	J3590	Not otherwise classified biologics
	J9999	Not otherwise classified anti-neoplastic
	C9399	Unclassified or biologics (hospital outpatient only)
Drug: NDC	50242-0245-01	1,200 mg pertuzumab, 600 mg trastuzumab, and 30,000 units hyaluronidase per 15 mL
	50242-0260-01	600 mg pertuzumab, 600 mg trastuzumab, and 20,000 units hyaluronidase per 10 mL
Administration procedures: CPT	96401	Chemotherapy administration, subcutaneous or intramuscular; non-hormonal anti-neoplastic
Billable units	For miscellaneous HCPCS codes, 1 billable unit is generally equal to 1 dose. Payers might have different preferences for billing for PHESGO. Check with your local payers for specific billing unit information.	

CPT=Current Procedural Terminology; HCPCS=Healthcare Common Procedure Coding System; ICD-10-CM=International Classification of Diseases, 10th Revision, Clinical Modification; NDC=National Drug Code.

These codes are not all-inclusive; appropriate codes can vary by patient, setting of care, and payer. Correct coding is the responsibility of the provider submitting the claim for the item or service. Please check with the payer to verify codes and special billing requirements. Genentech does not make any representation or guarantee concerning reimbursement or coverage for any service or item.

Many payers will not accept unspecified codes. If you use an unspecified code, please check with your payer.

Indications & Important Safety Information

Early Breast Cancer

PHESGO™ (pertuzumab, trastuzumab, and hyaluronidase-zzxf) is indicated for use in combination with chemotherapy for

- the neoadjuvant treatment of adult patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node-positive) as part of a complete treatment regimen for early breast cancer (EBC)
- the adjuvant treatment of adult patients with HER2-positive early breast cancer (EBC) at high risk of recurrence

Select patients for therapy based on an FDA-approved companion diagnostic test.

Metastatic Breast Cancer

PHESGO™ is indicated for use in combination with docetaxel for the treatment of adult patients with HER2-positive metastatic breast cancer (MBC) who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease.

Select patients for therapy based on an FDA-approved companion diagnostic test.

Please see full Prescribing Information and reverse for additional Important Safety Information, including BOXED WARNINGS.

Important Safety Information

BOXED WARNINGS: Cardiomyopathy, Embryo-Fetal Toxicity, and Pulmonary Toxicity

- **PHESGO administration can result in subclinical and clinical cardiac failure. The incidence and severity was highest in patients receiving PHESGO with anthracycline-containing chemotherapy regimens. Evaluate cardiac function prior to and during treatment with PHESGO. Discontinue PHESGO treatment in patients receiving adjuvant therapy and withhold PHESGO in patients with metastatic disease for clinically significant decrease in left ventricular function**
- **Exposure to PHESGO can result in embryo-fetal death and birth defects, including oligohydramnios and oligohydramnios sequence manifesting as pulmonary hypoplasia, skeletal abnormalities, and neonatal death. Advise patients of these risks and the need for effective contraception**
- **PHESGO administration can result in serious and fatal pulmonary toxicity. Discontinue PHESGO for anaphylaxis, angioedema, interstitial pneumonitis, or acute respiratory distress syndrome. Monitor patients until symptoms completely resolve**

Contraindications

PHESGO is contraindicated in patients with known hypersensitivity to pertuzumab, or trastuzumab, or hyaluronidase, or to any of its excipients.

Additional Important Safety Information

- Exacerbation of chemotherapy-induced neutropenia
- Hypersensitivity and administration-related reactions (ARRs): Monitor patients for systemic hypersensitivity reactions. Permanently discontinue PHESGO in patients who experience anaphylaxis or severe hypersensitivity reactions

Most Common Adverse Reactions

Early Breast Cancer

The most common adverse reactions (>30%) with PHESGO were alopecia, nausea, diarrhea, anemia, and asthenia.

Metastatic Breast Cancer (based on IV pertuzumab)

The most common adverse reactions (>30%) with pertuzumab in combination with trastuzumab and docetaxel were diarrhea, alopecia, neutropenia, nausea, fatigue, rash, and peripheral neuropathy.

You are encouraged to report side effects to Genentech and the FDA. You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at 1-888-835-2555.

Please see full Prescribing Information for additional Important Safety Information, including BOXED WARNINGS.

